arXiv:0806.2202v6 [math.NT] 23 Sep 2008

EXPLICIT CONSTRUCTIONS OF THE NON-ABELIAN $\mathbf{p^3}$-EXTENSIONS OVER $\mathbb{Q}$

OZ BEN-SHIMOL

Abstract. Let p be an odd prime. Let F/k be a cyclic extension of degree p and of characteristic different from p. The explicit constructions of the non-abelian p^3-extensions over k, are induced by certain elements in $F(\mu_p)^*$. In this paper we let $k = \mathbb{Q}$ and present sufficient conditions for these elements to be suitable for the constructions. Polynomials for the non-abelian groups of order 27 over $\mathbb{Q}$ are constructed.

1. Introduction

Let p be an odd prime. Let $F/\mathbb{Q}$ be a C_p-extension, $K = \mathbb{Q}(\zeta_p)$, $L = F(\zeta_p) = FK$, where ζ_p is a fixed p^{th} root of unity. Let $\bar{\sigma}$ be a generator for $\mathrm{Gal}(L/K)$. The restriction $\sigma = \bar{\sigma}|_F$ generates $\mathrm{Gal}(F/\mathbb{Q}) \cong \mathrm{Gal}(L/K)$. Similarly, let $\bar{\tau}$ be a generator for $\mathrm{Gal}(L/F)$. The restriction $\tau = \bar{\tau}|_K$ generates $\mathrm{Gal}(K/\mathbb{Q}) \cong \mathrm{Gal}(L/F)$. $L/\mathbb{Q}$ is cyclic of order $p(p-1)$, generated by $\bar{\sigma}\bar{\tau}$.

Let e be a primitive root modulo p. We have the multiplicative homomorphisms [4, p.130, Corollary 8.1.5]

$$\begin{cases} \Phi = \Phi_{L/F} : L^* \to L^* \\ \Phi_{L/F}(x) = x^{e^{p-2}} \bar{\tau} x^{e^{p-3}} \cdots \bar{\tau}^{p-2} x, \quad x \in L^*, \end{cases}$$

and its restriction $\Phi_{K/\mathbb{Q}} : K^* \to K^*$. Clearly, Φ commutes with the norm:

$$\mathrm{Nr}_{L/K} \Phi_{L/F} = \Phi_{K/\mathbb{Q}} \mathrm{Nr}_{L/K} .$$

Let $\mathbb{H}_{p^3}$ and $C_{p^2} \rtimes C_p$ be the non-abelian groups of order p^3. From the generic construction of the non-abelian p^3-extensions (over fields with characteristic different from p), it follows that the elements $x \in L^*$ with $\Phi(\mathrm{Nr}_{L/K}(x)) \notin L^{*p}$ induce $\mathbb{H}_{p^3}$-extensions over $\mathbb{Q}$, while the elements $x \in L^*$ with $\Phi(\zeta_p \mathrm{Nr}_{L/K}(x)) \notin L^{*p}$ induce $C_{p^2} \rtimes C_p$-extensions over $\mathbb{Q}$, cf. Theorem 4.1 and Theorem 4.2.

Arne Ledet, in [3],[4], and [5, pages 130-132], constructs explicit extensions and the corresponding polynomials over $\mathbb{Q}$ for these groups, when $p = 3$ or $p = 5$, and $L = \mathbb{Q}(\zeta_{p^2})$. He provides a sufficient condition for an element $x \in \mathcal{O}_L$ to satisfy the above requirements, namely: $\mathrm{Nr}_{L/\mathbb{Q}}(x) = q$ is an unramified prime in L (that is, $q \neq 3$, $q \neq 5$, respectively). His argument for the sufficiency relies on unique factorization in $\mathcal{O}_K$.

Our goal is to generalize this idea for every p and for every C_p-extension $F/\mathbb{Q}$ the construction starts with (in [4], the construction starts with a C_3-extension inside $\mathbb{Q}_9$). More precisely, given a C_p-extension $F/\mathbb{Q}$, we provide sufficient conditions for

---

1991 *Mathematics Subject Classification.* 12F12; 11R18.

Key words and phrases. Constructive Galois Theory, Heisenberg group, Explicit Embedding problem.

an element $x \in L^*$ to satisfy $\Phi(\mathrm{Nr}_{L/K}(x)) \notin L^{*p}$ for $\mathbb{H}_{p^3}$, and $\Phi(\zeta_p \,\mathrm{Nr}_{L/K}(x)) \notin L^{*p}$ for $C_{p^2} \rtimes C_p$. In section 4 we explain the explicit constructions for those two groups, and show that there are infinitely many elements $x \in L^*$, suitable for both constructions. In section 5, we exhibit other polynomials for $\mathbb{H}_{27}$ and $C_9 \rtimes C_3$ over $\mathbb{Q}$.

2. Notation and Definitions

- The number ring of a given number field N is denoted by $\mathcal{O}_N$.
- If E is a finite group and N/k is a Galois extension with Galois group $\mathrm{Gal}(N/k) \cong E$, we then say that N/K is an E-extension, or that N is an E-extension over k. Any polynomial f over k with splitting field N is called E-polynomial over k.
- A cyclic group of order m will denoted by C_m.
- As in the introduction: $F/\mathbb{Q}$ is a C_p-extension, $K = \mathbb{Q}(\zeta_p)$ and $L = FK$ – their compositum. Let $\bar{\sigma}$ be a fixed generator for $\mathrm{Gal}(L/K)$. Thus $\sigma = \bar{\sigma}|_F$ generates $\mathrm{Gal}(F/\mathbb{Q}) \cong \mathrm{Gal}(L/K)$. Let $\bar{\tau}$ be a fixed generator for $\mathrm{Gal}(L/F)$. Thus $\tau = \bar{\tau}|_K$ generates $\mathrm{Gal}(K/\mathbb{Q}) \cong \mathrm{Gal}(L/F)$.
- If e is a primitive root modulo p, then we have the homomorphism Φ defined in the introduction.
- Let $\mathcal{F}_L, \mathcal{F}_K$ be the multiplicative groups consisting of the fractional ideals of L,K, respectively. We have analogous homomorphisms:

$$\begin{cases} \widetilde{\Phi}_{L/F} : \mathcal{F}_L \to \mathcal{F}_L \\ \widetilde{\Phi}_{L/F}(I) = I^{e^{p-2}} \bar{\tau} I^{e^{p-3}} \cdots \bar{\tau}^{p-2} I, \quad I \in \mathcal{F}_L, \end{cases}$$

and its restriction

$$\begin{cases} \widetilde{\Phi} = \widetilde{\Phi}_{K/\mathbb{Q}} : \mathcal{F}_K \to \mathcal{F}_K \\ \widetilde{\Phi}_{K/\mathbb{Q}}(I) = \widetilde{\Phi}_{L/F}(\mathcal{O}_L I) \cap K, \quad I \in \mathcal{F}_K. \end{cases}$$

Clearly, $\mathrm{Nr}_{L/K} \widetilde{\Phi}_{L/F} = \widetilde{\Phi}_{K/\mathbb{Q}} \mathrm{Nr}_{L/K}$. [remark: let $I \in \mathcal{F}_L$, then $\mathrm{Nr}_{L/K}(I) = \mathcal{O}_K \cap I \bar{\sigma} I \cdots \bar{\sigma}^{p-1} I$ (see Marcus [6] or Ribenboim [8])].

- Let $x \in L^*$ with $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \pm q_1^{l_1} \cdots q_n^{l_n}$, $n > 0$, $l_i \in \mathbb{Z} - \{0\}$, and the q_i's are distinct rational primes. We denote by $J_i(x)$, $i = 1, \ldots n$, the fractional ideals of L which satisfy:

$$\mathcal{O}_L x = J_1(x) \cdots J_n(x) \quad \text{and} \quad \mathrm{Nr}_{L/\mathbb{Q}}(J_i(x)) = \mathbb{Z} q_i^{l_i}.$$

- For each i we set $I_i(x) := \mathrm{Nr}_{L/K}(J_i(x))$. Clearly, $\mathrm{Nr}_{K/\mathbb{Q}}(I_i(x)) = \mathbb{Z} q_i^{l_i}$.
- If for some i, q_i splits completely in K and $I_i(x) = P_1^{\beta_1} P_2^{\beta_2} \cdots P_{p-1}^{\beta_{p-1}}$, where the P_j's are the distinct prime ideals in K lying above q_i, then we define $\chi(I_i(x))$ to be the integer

$$\chi(I_i(x)) = e^{p-2}\beta_1 + e^{p-3}\beta_{p-1} + \ldots e\beta_3 + \beta_2.$$

3. The Main Results

Theorem 3.1. *Let* $x \in L^*$, $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \pm q_1^{l_1} \cdots q_n^{l_n}$, $n \geq 0$, $l_i \in \mathbb{Z} - \{0\}$, q_i *a prime. Then* $\Phi(\mathrm{Nr}_{L/K}(x))$ *does not generate a* p^{th} *power (fractional) ideal of* L*, if and only if* $n > 0$ *and there exists an* i *such that*

(a) q_i *splits completely in* L.

(b) p *does not divide* $\chi(I_i(x))$.

Proof. We assume $n > 0$. Denote $\gamma = \mathrm{Nr}_{L/K}(x)$. Clearly, $\mathcal{O}_K\gamma = I_1(x)\cdots I_n(x)$. No prime factor of $I_i(x)$ is conjugate to any prime factor of $I_j(x)$ $(i \neq j)$. It follows that $\mathcal{O}_L\Phi(\gamma) \in \mathcal{F}_L^p$ if and only if $\mathcal{O}_L\widetilde{\Phi}(I_i(x)) \in \mathcal{F}_L^p$ for every $i = 1, \ldots, n$.

We fix some i and set $q = q_i$, $l = l_i$, $J = J_i(x)$, $I = I_i(x)$. Thus, J is a fractional ideal of L, $I = \mathrm{Nr}_{L/K}(J)$, and $\mathrm{Nr}_{L/\mathbb{Q}}(J) = \mathrm{Nr}_{K/\mathbb{Q}}(I) = \mathbb{Z}q^l$.

We separate the rest of the proof into two cases.

Case 1: q does not split completely in L

Let g be the decomposition number of q in $K/\mathbb{Q}$. Clearly, $\tau^g I = I$. Now,

$$\widetilde{\Phi}(I) = \prod_{j=0}^{p-2} \tau^j I^{e^{p-(j+2)}} = \prod_{j=1}^{\frac{p-1}{g}} \prod_{i=0}^{g-1} \tau^i I^{e^{p-((j-1)g+i+2)}} = \prod_{i=0}^{g-1} \tau^i I^{S_g(i)},$$

where,

$$S_g(i) = \sum_{j=1}^{\frac{p-1}{g}} e^{p-((j-1)g+i+2)} = e^{p-(i+2)} \cdot \frac{e^{-(p-1)}-1}{e^{-g}-1}.$$

We see that if q does not split completely in K then $S_g(i) \equiv 0 \mod p$ for all $i = 0, \ldots g-1$. Therefore, if q does not split completely in K then $\widetilde{\Phi}(I) \in \mathcal{F}_K^p$.

Suppose that q splits completely in K. If q is ramified in F then $q \equiv 1 \mod p$ and any prime ideal P in K lying above q is then totally ramified in L, equivalently, $\mathcal{O}_L P \in \mathcal{F}_L^p$, hence $\mathcal{O}_L I \in \mathcal{F}_L^p$..

Finally, suppose that q is inert in F. Then any prime in K lying above q is inert in L. Hence, $\bar{\sigma}J = J$, thus $\mathcal{O}_L I = J^p \in \mathcal{F}_L^p$.

We conclude that if q does not split completely in L, then $\mathcal{O}_L\widetilde{\Phi}(I) \in \mathcal{F}_L^p$.

Case 2: q splits completely in L

Suppose that q splits completely in L, and let $\mathcal{O}_K q = P_1 \ldots P_{p-1}$ be the prime decomposition of q in K. τ, as a permutation acting on the $(p-1)$-set consisting of the P_j's, is the $(p-1)$-cycle $(P_1, \ldots, P_{p-1})$ (say). We shall denote by $\check{\tau}$ the corresponding $(p-1)$-cycle in the symmetric group; $\check{\tau} = (1, \ldots, p-1)$. Now, if

$$I = P_1^{\beta_1} P_2^{\beta_2} \cdots P_{p-1}^{\beta_{p-1}},$$

then

$$\widetilde{\Phi}(I) = \prod_{j=1}^{p-1} \Phi(P_j)^{\beta_j} = \prod_{j=1}^{p-1}\prod_{k=0}^{p-2} \bar{\tau}^k P_j^{e^{(p-(k+2))}\beta_j} = \prod_{j=1}^{p-1} P_j^{\chi_j(I)}, \tag{1}$$

where,

$$\chi_j(I) = \sum_{k=0}^{p-2} e^{(p-(k+2))}\beta_{\check{\tau}^{-k}(j)}, \quad j = 1, \ldots, p-1.$$

Clearly, the primes P_j split completely in L, hence, $\mathcal{O}_L\widetilde{\Phi}(I) \in \mathcal{F}_L^p$ if and only if $\widetilde{\Phi}(I) \in \mathcal{F}_K^p$. Also, $\widetilde{\Phi}(I) \in \mathcal{F}_K^p$ if and only if $\chi_j(I) \equiv 0 \mod p$ for all $j = 1, \ldots, p-1$. Note that $\chi_{j+1}(I) \equiv e \cdot \chi_j(I) \mod p$ for all j. Therefore, if p divides some $\chi_j(I)$, then p must divides $\chi_1(I), \ldots, \chi_{p-1}(I)$. Since we consider the $\chi_j(I)$'s modulo p, we take $j = 1$ and write $\chi(I)$ instead of $\chi_1(I)$. We conclude that $\mathcal{O}_L\widetilde{\Phi}(I) \in \mathcal{F}_L^p$ if and only if p divides $\chi(I)$, as required.

If $n = 0$ then the assertion is clear. $\square$

Corollary 3.2. *Let* $x \in L^*$, $\operatorname{Nr}_{L/\mathbb{Q}}(x) = \pm q_1^{l_1} \cdots q_n^{l_n}$, $n \geq 1$, $l_i \in \mathbb{Z} - \{0\}$, q_i *a prime. Suppose that there exists an* i *such that* q_i *splits completely in* L *and* p *does not divide* $\chi(I_i(x))$*. Then* $\Phi(\operatorname{Nr}_{L/K}(x)), \Phi(\zeta_p \operatorname{Nr}_{L/K}(x)) \notin L^{*p}$. □

Let $x \in L^*$ which satisfies condition (a) of Theorem 3.1. Let q_i be a prime which splits completely in L with $\operatorname{Nr}_{K/\mathbb{Q}}(I_i(x)) = \mathbb{Z}q_i^{l_i}$. If P is a prime ideal in K lying above q_i then $\operatorname{Nr}_{K/\mathbb{Q}}(P) = \mathbb{Z}q_i$ (since q_i splits completely in K). Therefore, if $I_i(x) = P_1^{\beta_1} P_2^{\beta_2} \cdots P_{p-1}^{\beta_{p-1}}$, where the P_j's are the distinct prime ideals in K lying above q_i, then

$$\mathbb{Z}q_i^{l_i} = \operatorname{Nr}_{K/\mathbb{Q}}(I_i(x)) = \prod_{j=1}^{p-1} \operatorname{Nr}_{K/\mathbb{Q}} P_j^{\beta_j} = \mathbb{Z}q_i^{\beta_1 + \ldots \beta_{p-1}}.$$

Thus,

$$\beta_1 + \ldots + \beta_{p-1} = l_i. \tag{2}$$

It follows immediately that if $I_i(x)$ is of the form $I_i(x) = P^l$ (for some prime ideal P in K lying above q_i and for some $l \in \mathbb{Z} - \{0\}$), then p divides $\chi(I_i(x))$ if and only if p divides l. We have

Corollary 3.3. *Let* $x \in L^*$, $\operatorname{Nr}_{L/\mathbb{Q}}(x) = \pm q_1^{l_1} \cdots q_n^{l_n}$, $n \geq 1$, $l_i \in \mathbb{Z} - \{0\}$, q_i *a prime. Suppose that there exists an* i *such that* q_i *splits completely in* L *and* $I_i(x) = P^{l_i}$*, where* P *is a prime ideal of* $\mathcal{O}_K$*,* p *does not divide* l_i*. Then* $\Phi(\operatorname{Nr}_{L/K}(x)), \Phi(\zeta_p \operatorname{Nr}_{L/K}(x)) \notin L^{*p}$. □

Let $\mathcal{H}(L)$ be the Hilbert class field over L (see [6, Chapter 8]), and let q be a rational prime which splits completely in $\mathcal{H}(L)$. Then any prime ideal of L lying above q is principal in L.

Corollary 3.4. *There are infinitely many elements* $x \in L^*$ *such that*
1. $\Phi(\operatorname{Nr}_{L/K}(x)) \notin L^{*p}$*, and*
2. $\Phi(\zeta_p \operatorname{Nr}_{L/K}(x)) \notin L^{*p}$.

Proof. By Chebotarev's Density Theorem, there are infinitely many rational primes which split completely in $\mathcal{H}(L)$. If q is such a prime, let Q be a prime ideal in L lying above q. So $Q = \mathcal{O}_L x$ for some $x \in \mathcal{O}_L$. Clearly, $\operatorname{Nr}_{L/\mathbb{Q}}(x) = q$. The assertion then follows from Corollary 3.3. □

3.1. Examples.

We shall illustrate the above results on C_p-extensions $F/\mathbb{Q}$ of type (S_1) [10, Chap. 2]. For that purpose we prove the following lemma.

Lemma 3.5. *Let* r *be a rational prime,* $r \equiv 1 \mod k$, $k \in \mathbb{N}$*. Let* $m = m(r)$ *be a primitive root modulo* r*. Consider the sum*

$$\delta_k(r) = \zeta_r + \zeta_r^{m^k} + \zeta_r^{m^{2k}} + \ldots + \zeta_r^{m^{\left(\frac{r-1}{k}-1\right)k}}. \tag{3}$$

Then $\mathbb{Q}(\delta_k(r))/\mathbb{Q}$ *is a* C_k*-extension.*

Proof. $\operatorname{Gal}(\mathbb{Q}(\zeta_r)/\mathbb{Q})$ is cyclic of order $r-1$, generated by the automorphism $\sigma : \zeta_r \mapsto \zeta_r^m$. By the Fundamental Theorem of Galois Theory, it is enough to prove $\mathbb{Q}(\delta_k(r)) = \mathbb{Q}(\zeta_r)^{\sigma^k}$. The inclusion $\mathbb{Q}(\delta_k(r)) \subseteq \mathbb{Q}(\zeta_r)^{\sigma^k}$ is because σ^k moves cyclically the summands of (3) (an alternative argument: $\delta_k(r)$ is the image of ζ_r under the trace map $\operatorname{Tr}_{\mathbb{Q}(\zeta_r)/\mathbb{Q}(\zeta_r)^{\sigma^k}}$). Suppose that $\mathbb{Q}(\delta_k(r)) \subsetneqq \mathbb{Q}(\zeta_r)^{\sigma^k}$. There exist a proper divisor d of k such that $\mathbb{Q}(\delta_k(r)) = \mathbb{Q}(\zeta_r)^{\sigma^d}$. In particular, $\sigma^d(\delta_k(r)) = \delta_k(r)$, or

$$\sum_{j=0}^{\frac{r-1}{k}-1} \zeta_r^{m^{jk+d}} - \sum_{j=0}^{\frac{r-1}{k}-1} \zeta_r^{m^{jk}} = 0. \tag{4}$$

The summands in (4) are distinct. For if $\zeta_r^{m^{jk+d}} = \zeta_r^{m^{ik}}$ for some $i, j = 0, 1, \ldots, \frac{r-1}{k} - 1$, $j \geq i$, then $m^{(j-i)k+d} \equiv 1(\mod r)$. m is primitive modulo r so $r-1$ divides $(j-i)k+d$. But, $(j-i)k+d < (\frac{r-1}{k}-1)k+k = r-1$, a contradiction. To this end, there are $2(r-1)/k$ $(\leq r-1)$ summands, and dividing each of them by ζ_r gives us a linear dependence among $1, \zeta_r, \zeta_r^2, \ldots, \zeta_r^{r-2}$, a contradiction. $\mathbb{Q}(\delta_k(r)) = \mathbb{Q}(\zeta_r)^{\sigma^k}$, as required. □

Example 3.6. $p = 3$, $r = 7$, $m(r) = 3$. $\delta_3(7) = \zeta_7 + \zeta_7^{-1}$. $F/\mathbb{Q} = \mathbb{Q}(\delta_3(7))/\mathbb{Q}$. Thus, $L/K = \mathbb{Q}(\zeta_3, \delta_3(7))/\mathbb{Q}(\zeta_3)$ is a C_3-extension, generated by $\bar{\sigma} : \zeta_7 \mapsto \zeta_7^2$, $\zeta_3 \mapsto \zeta_3$. Let $x = \delta_3(7) + \zeta_3$. $\gamma = \mathrm{Nr}_{L/K}(x) = 3 - \zeta_3$, $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \mathrm{Nr}_{K/\mathbb{Q}}(\gamma) = 13$ $(\tau : \zeta_3 \mapsto \zeta_3^{-1})$. 13 splits completely in L. By Corollary 3.3, $\Phi(\gamma)$, $\zeta_3\Phi(\gamma) \notin L^{*3}$.

Example 3.7. $p = 3$, $r = 19$, $m(r) = 2$.

$$\delta_3(19) = \zeta_{19} + \zeta_{19}^{-1} + \zeta_{19}^{7} + \zeta_{19}^{-7} + \zeta_{19}^{8} + \zeta_{19}^{-8}.$$

$F/\mathbb{Q} = \mathbb{Q}(\delta_3(19))/\mathbb{Q}$. Thus, $L/K = \mathbb{Q}(\zeta_3, \delta_3(19))/\mathbb{Q}(\zeta_3)$ is a C_3-extension, generated by $\bar{\sigma} : \zeta_{19} \mapsto \zeta_{19}^6$, $\zeta_3 \mapsto \zeta_3$. Let $x = \delta_3(19) + \zeta_3 + 1$. $\gamma = \mathrm{Nr}_{L/K}(x) = -7\zeta_3$, $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \mathrm{Nr}_{K/\mathbb{Q}}(\gamma) = 7^2$ and 7 splits completely in L. However, $\Phi(\gamma)$ generates a third power ideal in $\mathcal{O}_K$ although $\Phi(\gamma)$ and $\zeta_3\Phi(\gamma)$ are not third power elements in L^*. Therefore, Theorem 3.1, which is an *ideal-theoretic* criterion, is failed while testing this x. On the other hand, it tells us something about the prime factorization of $\mathcal{O}_L x$ in L: Since x satisfies condition (a) of this theorem and does not satisfy condition (b), we have

$$\chi(I_1(x)) = 2\beta_1 + \beta_2 \equiv 0 \mod 3.$$

Also, by equation (2) we have $\beta_1 + \beta_2 = 2$. Hence, $\beta_1 = \beta_2 = 1$, thus $I = \mathcal{O}_K 7$. $\mathcal{O}_L x$ is therefore a product of two distinct primes of O_L, each of them has norm 7 over $\mathbb{Q}$.

Example 3.8. $p = 3$, $r = 73$, $m(r) = 5$. (Remark: $\mathcal{O}_K = \mathbb{Z}[\zeta_{73}]$ is not a unique factorization domain). $F/\mathbb{Q} = \mathbb{Q}(\delta_3(73))/\mathbb{Q}$. Thus, $L/K = \mathbb{Q}(\zeta_3, \delta_3(73))/\mathbb{Q}(\zeta_3)$ is a C_3-extension, generated by $\bar{\sigma} : \zeta_{73} \mapsto \zeta_{73}^{24}$, $\zeta_3 \mapsto \zeta_3$. Let $x = \delta_3(73) - \zeta_3 + 1$. $\gamma = \mathrm{Nr}_{L/K}(x) = 21\zeta_3$, $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \mathrm{Nr}_{K/\mathbb{Q}}(\gamma) = 3^2 7^2$. Thus, $\Phi(\gamma) = 21^3\zeta_3 \notin L^{*3}$.

Example 3.9. $p = 5, r = 11, m(r) = 2$. $\delta_5(11) = \zeta_{11} + \zeta_{11}^{-1}$. $F/\mathbb{Q} = \mathbb{Q}(\delta_5(11))/\mathbb{Q}$. Thus, $L/K = \mathbb{Q}(\zeta_5, \delta_5(11))/\mathbb{Q}(\zeta_5)$ is a C_5-extension, generated by $\bar{\sigma} : \zeta_{11} \mapsto \zeta_{11}^2$, $\zeta_5 \mapsto \zeta_5$. Let $x = \delta_5(11) - \zeta_5$. $\gamma = \mathrm{Nr}_{L/K}(x)$, $\mathrm{Nr}_{L/\mathbb{Q}}(x) = \mathrm{Nr}_{K/\mathbb{Q}}(\gamma) = 991$ (991 is a rational prime). 991 splits completely in L, hence $\Phi(\gamma)$ does not generate a fifth power ideal of $\mathcal{O}_L$.

4. Explicit Constructions of the non-Abelian p^3-Extensions

4.1. Let N/k be a Galois extension with Galois group $G = \mathrm{Gal}(N/k)$. Let $\pi : E \twoheadrightarrow G$ be an epimorphism of a given finite group E onto G. *The Galois theoretical embedding problem* is to find a pair $(T/k, \varphi)$ consisting of a Galois extension T/k which contains N/k, and an isomorphism $\varphi : \mathrm{Gal}(T/k) \to E$ such that $\pi \circ \varphi =$

$\mathrm{Res}_{T/N}$, where $\mathrm{Res}_{T/N} : \mathrm{Gal}(T/k) \twoheadrightarrow G$ is the restriction map. We say that the embedding problem is *given by* N/k *and* $\pi : E \twoheadrightarrow G$. If such a pair $(T/k, \varphi)$ does exist, we say that the embedding problem is *solvable.* T is called a *solution field*, and φ is called a *solution isomorphism.* $A := \ker \pi$ is called the *kernel of the embedding problem.* If A is cyclic of order m, the characteristic of k does not divide m, and N^* contains μ_m (the multiplicative group consisting of the m^{th} roots of unity), then the embedding problem is of *Brauer type* (see [4]).

If φ is required only to be a monomorphism, we say that the embedding problem is *weakly solvable.* In that case, the pair $(T/k, \varphi)$ is called an *improper solution* (or a *weak solution*). An improper solution can also be interpret in terms of Galois algebra ([1], [3] Chapter 4). The monograph [2] treats the embedding problem from this point of view. In the cases discussed in this paper, any improper solution will automatically be a solution; the kernel A will be contained in the Frattini subgroup of E (see [2], Chapter.6, Corollary 5).

The constructive approach to embedding problems is to describe explicitly the solution field T (in case it exists). We shall consider a series of embedding problems, in each T/N will be a C_p-extension, and the goal is to exhibit an explicit primitive element of T over N (See also [7] for further details). In practice, given a set of generators $\gamma_1, \ldots, \gamma_d$ for G, we must extend them to k-automorphisms $\bar{\gamma}_1, \ldots, \bar{\gamma}_d$ of a larger field T, with common fixed field k. T/k is then a Galois extension generated by $\bar{\gamma}_1, \ldots, \bar{\gamma}_d, \bar{\gamma}_{d+1}, \ldots, \bar{\gamma}_c$, where $\bar{\gamma}_{d+1}, \ldots, \bar{\gamma}_c$ generate $\mathrm{Gal}(T/N)$. The extensions should be constructed in such a way that the $\bar{\gamma}_i$'s behave as a given set of c generators for E. T is then a solution field, and the isomorphism which takes each $\bar{\gamma}_i$ to the corresponding generator of E is a solution isomorphism.

4.2. Let N be a field of characteristic different from p. $\mu_p \subseteq N^*$, and let $a \in N^* \setminus N^{*p}$. Then $N(\sqrt[p]{a})/N$ is a C_p-extension. A generator for $\mathrm{Gal}(N(\sqrt[p]{a})/N)$ is given by $\sqrt[p]{a} \mapsto \zeta_p \sqrt[p]{a}$. The only elements $\theta \in N(\sqrt[p]{a})$ with the property: $\theta^p \in N$, are those of the form $z(\sqrt[p]{a})^j$ for some $z \in N$, $j = 0, 1, \ldots, p-1$.

Suppose further that N/k is a Galois extension, then $N(\sqrt[p]{a})/k$ is a Galois extension if and only if, for each $\gamma \in \mathrm{Gal}(N/k)$, there exist $i_\gamma \in \mathbb{Z} \setminus p\mathbb{Z}$ such that $\gamma a / a^{i_\gamma} \in N^{*p}$.

4.3. Let E be a non-abelian group of order p^3. Up to isomorphism, E is necessarily one of the following two groups:

1. The Heisenberg group:

$$\mathbb{H}_{p^3} = \langle\ u, v, w\ :\ u^p = v^p = w^p = 1,\ wu = uw,\ wv = vw,\ vu = uvw\ \rangle.$$

In other words, $\mathbb{H}_{p^3}$ is generated by two elements u, v of order p, such that their commutator w is central. It can be realized as the subgroup of $\mathrm{GL}_3(\mathbb{Z}/\mathbb{Z}p)$ consisting of upper triangular matrices with 1's in the diagonal.

2. The semidirect product:

$$C_{p^2} \rtimes C_p = \langle\ u, v\ :\ u^{p^2} = v^p = 1,\ vu = u^{p+1}v\ \rangle.$$

For the classification of the non-abelian groups of order p^3, see [9, page 67].

4.4. The aim of the following Theorem 4.1 (resp. Theorem 4.2) is to describe *how* $\mathbb{H}_{p^3}$-extensions (resp. $C_{p^2} \rtimes C_p$-extensions) are constructed over $\mathbb{Q}$, starting with elements $x \in L^*$ with $\Phi(\mathrm{Nr}_{L/K}(x)) \notin L^{*p}$ (resp. $\Phi(\zeta_p \mathrm{Nr}_{L/K}(x)) \notin L^{*p}$). These constructions are essentially known - they can be obtained as private cases of [4, Theorem 2.4.1 and Corollary 8.1.5]. We give a direct and self contained proof, consistent with our notation and with subsection 4.1.

Theorem 4.1 (The case $E = \mathbb{H}_{p^3}$).
Let $x \in L^*$ *such that* $b = b(x) = \Phi(\mathrm{Nr}_{L/K}(x)) \notin L^{*p}$. *Then* $G = \mathrm{Gal}(L(\sqrt[p]{b})/K) \cong C_p \times C_p$, *generated by* $\bar{\bar{\sigma}}$ *and* η, *where,* $\bar{\bar{\sigma}} : \sqrt[p]{b} \mapsto \sqrt[p]{b}$, $\eta : \sqrt[p]{b} \mapsto \zeta_p \sqrt[p]{b}$, $\bar{\bar{\sigma}}|_L = \bar{\sigma}$, $\eta|_L = \mathrm{id}_L$. *Moreover, the (Brauer type) embedding problem given by* $L(\sqrt[p]{b})/K$ *and* $\pi : E \twoheadrightarrow G$ *is solvable, where* $\pi : u \mapsto \bar{\bar{\sigma}}$, $v \mapsto \eta$; *a solution field is* $M = L(\sqrt[p]{\omega}, \sqrt[p]{b})$, $\omega = \Phi(\beta)$, $\beta = x^{p-1}\bar{\sigma}x^{p-2} \ldots \bar{\sigma}^{p-2}x$.

Since $\omega \in \Phi(L^*)$, $M/\mathbb{Q}$ *is a Galois extension with* $\mathrm{Gal}(M^\kappa/\mathbb{Q}) \cong E$, *where* κ *is the extension of* $\bar{\tau}$ *to* M, *given by* $\kappa : \sqrt[p]{\omega} \mapsto \beta^{(1-e^{p-1})/p}\left(\sqrt[p]{\omega}\right)^e$, *and* $M^\kappa = F(\alpha)$, $\alpha = \sqrt[p]{\omega} + \kappa\sqrt[p]{\omega} + \ldots + \kappa^{p-2}\sqrt[p]{\omega}$. *Finally,* $F(\alpha)$ *is the splitting field of the minimal polynomial for* α *over* $\mathbb{Q}$.



Proof. Since $\bar{\sigma}b = b$, $L(\sqrt[p]{b})/K$ is a Galois extension. We extend $\bar{\sigma}$ to $\mathrm{Gal}(L(\sqrt[p]{b})/K)$ by $\bar{\bar{\sigma}}(\sqrt[p]{b}) = \sqrt[p]{b}$. η generates $\mathrm{Gal}(L(\sqrt[p]{b})/L)$, and $\bar{\sigma}$ generates $\mathrm{Gal}(L/K)$. It follows that $\mathrm{Gal}(L(\sqrt[p]{b})/K)$ is generated by $\bar{\bar{\sigma}}$ and η. Clearly, $\bar{\bar{\sigma}}\eta = \eta\bar{\bar{\sigma}}$ and $\bar{\bar{\sigma}}^p = \eta^p = \mathrm{id}_{L(\sqrt[p]{b})}$, thus, $L(\sqrt[p]{b})/K$ is a $C_p \times C_p$-extension $[L(\sqrt[p]{b})$ is a Kummer extension of K of exponent $p]$.

$\eta(\omega) = \omega$ and,

$$\frac{\bar{\bar{\sigma}}(\omega)}{\omega} = \frac{\bar{\sigma}(\Phi(\beta))}{\Phi(\beta)} = \frac{\Phi(\mathrm{Nr}_{L/K}(x))}{\Phi(x)^p} = \left(\frac{\sqrt[p]{b}}{\Phi(x)}\right)^p \in L(\sqrt[p]{b})^{*^p}, \tag{5}$$

thus, M/K is a Galois extension. We extend $\bar{\bar{\sigma}}$ and η to $\tilde{\sigma}, \tilde{\eta} \in \mathrm{Gal}(M/K)$ by

$$\tilde{\sigma}(\sqrt[p]{\omega}) = \frac{\sqrt[p]{b}}{\Phi(x)}\sqrt[p]{\omega}, \qquad \tilde{\eta}(\sqrt[p]{\omega}) = \zeta_p\sqrt[p]{\omega}, \tag{6}$$

respectively. Note that $M/L(\sqrt[p]{b})$ is a C_p-extension. For if $\omega \in L(\sqrt[p]{b})^{*^p}$ then $\omega = z^p b^j$ for some $z \in L$, $j = 0, 1, \ldots p-1$. From (5) it follows that $b = (\Phi(x)(\bar{\sigma}(z)/z))^p \in L^{*p}$ - a contradiction.

$\mathrm{Gal}(M/K)$ is then generated by $\tilde{\sigma}, \tilde{\eta}$ and $\tilde{\lambda}$, where $\tilde{\lambda}(\sqrt[p]{\omega}) = \zeta_p\sqrt[p]{\omega}$ - a generator for $\mathrm{Gal}(M/L(\sqrt[p]{b}))$. Clearly, $\tilde{\eta}$ and $\tilde{\lambda}$ is of order p, and since

$$\tilde{\sigma}^p(\sqrt[p]{\omega}) = \frac{b}{\Phi(x)\bar{\sigma}(\Phi(x))\cdots\bar{\sigma}^{p-1}(\Phi(x))}\sqrt[p]{\omega} = \frac{b}{\mathrm{Nr}_{L/K}(\Phi(x))}\sqrt[p]{\omega} = \frac{b}{\Phi(\mathrm{Nr}_{L/K}(x))}\sqrt[p]{\omega} = \sqrt[p]{\omega},$$

$\tilde{\sigma}$ is of order p. $\tilde{\lambda}$ commutes with both $\tilde{\sigma}$ and $\tilde{\eta}$, and the relation $\tilde{\eta}\tilde{\sigma} = \tilde{\sigma}\tilde{\eta}\tilde{\lambda}$ holds. Therefore, the group isomorphism $\varphi : \mathrm{Gal}(M/K) \cong \mathbb{H}_{p^3}$, sending $\tilde{\sigma} \mapsto u$, $\tilde{\eta} \mapsto v$, $\tilde{\lambda} \mapsto w$, is a solution isomorphism for the given embedding problem.

Now,

$$\frac{\bar{\tau}\Phi(x)}{\Phi(x)^e} = \Phi\left(\frac{\bar{\tau}x}{x^e}\right) = \frac{\bar{\tau}x^{e^{p-2}}}{x^{e^{p-1}}} \cdot \frac{\bar{\tau}^2 x^{e^{p-3}}}{\bar{\tau}x^{e^{p-2}}} \cdots \frac{\bar{\tau}^{p-2}x^e}{\bar{\tau}^{p-3}x^{e^2}} \cdot \frac{x}{\bar{\tau}^{p-2}x^e} = x^{1-e^{p-1}},$$

hence,

$$\frac{\bar{\tau}\omega}{\omega^e} = \frac{\bar{\tau}\Phi\left(x^{p-1}\bar{\sigma}x^{p-2}\cdots\bar{\sigma}^{p-2}x\right)}{\Phi\left(x^{p-1}\bar{\sigma}x^{p-2}\cdots\bar{\sigma}^{p-2}x\right)^e} = \left(\frac{\bar{\tau}\Phi(x)}{\Phi(x)^e}\right)^{p-1} \bar{\sigma}\left(\frac{\bar{\tau}\Phi(x)}{\Phi(x)^e}\right)^{p-2} \cdots \bar{\sigma}^{p-2}\left(\frac{\bar{\tau}\Phi(x)}{\Phi(x)^e}\right)$$

$$= (x^{p-1}\bar{\sigma}x^{p-2}\cdots\bar{\sigma}^{p-2}x)^{1-e^{p-1}} = \left(\beta^{\frac{1-e^{p-1}}{p}}\right)^p \in L^{*p},$$

and

$$\frac{\bar{\tau}b}{b^e} = \frac{\bar{\tau}\Phi(\mathrm{Nr}_{L/K}(x))}{\Phi(\mathrm{Nr}_{L/K}(x))^e} = \mathrm{Nr}_{L/K}\left(\frac{\bar{\tau}\Phi(x)}{\Phi(x)^e}\right) = \left(\mathrm{Nr}_{L/K}(x)^{\frac{1-e^{p-1}}{p}}\right)^p \in L^{*p}.$$

Thus, $M/\mathbb{Q}$ is a Galois extension. We extend $\bar{\tau}$ to $\kappa \in \mathrm{Gal}(M/\mathbb{Q})$ by

$$\kappa(\sqrt[p]{\omega}) = \beta^{(1-e^{p-1})/p}\left(\sqrt[p]{\omega}\right)^e, \qquad \kappa(\sqrt[p]{b}) = \mathrm{Nr}_{L/K}(x)^{(1-e^{p-1})/p}(\sqrt[p]{b})^e.$$

Since κ is an extension of τ to M, $\mathrm{Gal}(M/\mathbb{Q})$ is generated by $\tilde{\sigma}$,$\tilde{\eta}$,$\tilde{\lambda}$, and κ. It is easy to verify that κ is central and has order $p-1$ [while doing it, we take $\tau : \zeta_p \mapsto \zeta_p^e$. Also, note that κ is the unique extension of $\bar{\tau}$ of order $p-1$]. It follows that $\mathrm{Gal}(M/\mathbb{Q}) \cong \mathbb{H}_{p^3} \times C_{p-1}$ and $M^\kappa/\mathbb{Q}$ is Galois with $\mathrm{Gal}(M^\kappa/\mathbb{Q}) \cong \mathbb{H}_{p^3}$, as required.

Clearly, $F(\alpha) \subseteq M^\kappa$. We have the short exact sequence

$$\begin{tikzcd} 1 \arrow[r] & \mathrm{Gal}(M/F) \arrow[r, "\imath"] & \mathrm{Gal}(M/\mathbb{Q}) \arrow[r, "\mathrm{Res}"] & \mathrm{Gal}(F/\mathbb{Q}) \arrow[r] & 1, \end{tikzcd}$$

It follows that $\mathrm{Gal}(M/F)$ is generated by $\tilde{\eta}$,$\tilde{\lambda}$, and κ. Thus, $F(\alpha) = M^\kappa$ if and only if $\tilde{\eta}\alpha \neq \alpha$ and $\tilde{\lambda}\alpha \neq \alpha$. But $\tilde{\eta}\alpha = \alpha$ or $\tilde{\lambda}\alpha = \alpha$ imply linear dependence among the elements $\sqrt[p]{\omega}$, $\sqrt[p]{\omega}^e, \ldots, \sqrt[p]{\omega}^{e^{p-2}}$ over L, a contradiction.

To this end, $\mathrm{Gal}(F(\alpha)/\mathbb{Q}(\alpha)) \cong \mathrm{Gal}(F/\mathbb{Q}) \cong C_p$, and $\mathbb{Q}(\alpha)/\mathbb{Q}$ is not a normal extension. Hence, $F(\alpha)$ is the splitting field for $\mathrm{Irr}(\alpha; \mathbb{Q})$. $\square$

Theorem 4.2 (The case $E = C_{p^2} \rtimes C_p$).
Let $x \in L^*$ such that $b = b(x) = \Phi(\zeta_p \mathrm{Nr}_{L/K}(x)) \notin L^{*p}$. Then $G = \mathrm{Gal}(L(\sqrt[p]{b})/K) \cong C_p \times C_p$, generated by $\bar{\bar{\sigma}}$ and η, where, $\bar{\bar{\sigma}} : \sqrt[p]{b} \mapsto \sqrt[p]{b}$, $\eta : \sqrt[p]{b} \mapsto \zeta_p\sqrt[p]{b}$, $\bar{\bar{\sigma}}|_L = \bar{\sigma}$, $\eta|_L = \mathrm{id}_L$. Moreover, the (Brauer type) embedding problem given by $L(\sqrt[p]{b})/K$ and $\pi : E \twoheadrightarrow G$ is solvable, where $\pi : u \mapsto \bar{\bar{\sigma}}$, $v \mapsto \eta$; a solution field is $M = L(\sqrt[p]{\omega}, \sqrt[p]{b})$, $\omega = \Phi(\beta\sqrt[p]{a})$, $\beta = x^{p-1}\bar{\sigma}x^{p-2}\ldots\bar{\sigma}^{p-2}x$, $L = K(\sqrt[p]{a})$.

Since $\omega \in \Phi(L^*)$, $M/\mathbb{Q}$ is a Galois extension with $\mathrm{Gal}(M^\kappa/\mathbb{Q}) \cong E$, where κ is the extension of $\bar{\tau}$ to M, given by $\kappa : \sqrt[p]{\omega} \mapsto (\beta\sqrt[p]{a})^{(1-e^{p-1})/p}(\sqrt[p]{\omega})^e$, and $M^\kappa = F(\alpha)$, $\alpha = \sqrt[p]{\omega} + \kappa\sqrt[p]{\omega} + \ldots + \kappa^{p-2}\sqrt[p]{\omega}$. Finally, $F(\alpha)$ is the splitting field of the minimal polynomial for α over $\mathbb{Q}$.

Proof. A few changes should be made in the proof of Theorem 4.1.
We can assume $\bar{\sigma} : \sqrt[p]{a} \mapsto \zeta_p \sqrt[p]{a}$. Next, $\eta\omega = \omega$ and,

$$\frac{\bar{\bar{\sigma}}(\omega)}{\omega} = \frac{\bar{\sigma}(\Phi(\beta))}{\Phi(\beta)} \cdot \Phi\left(\frac{\bar{\sigma}\sqrt[p]{a}}{\sqrt[p]{a}}\right) = \frac{\Phi(\zeta_p \operatorname{Nr}_{L/K}(x))}{\Phi(x)^p} = \left(\frac{\sqrt[p]{b}}{\Phi(x)}\right)^p \in L(\sqrt[p]{b})^{*p},$$

thus, M/K is a Galois extension. We extend $\bar{\bar{\sigma}}$ and η to $\tilde{\sigma}, \tilde{\eta} \in \operatorname{Gal}(M/K)$ as in [6]. $\operatorname{Gal}(M/K)$ is generated by $\tilde{\sigma}, \tilde{\eta}$ and $\tilde{\lambda}$, where $\tilde{\lambda}(\sqrt[p]{\omega}) = \zeta_p \sqrt[p]{\omega}$ - a generator for $\operatorname{Gal}(M/L(\sqrt[p]{b}))$. Clearly, $\tilde{\eta}$ is of order p. Now,

$$\tilde{\sigma}^p(\sqrt[p]{\omega}) = \frac{b}{\Phi(x)\bar{\sigma}(\Phi(x))\cdots\bar{\sigma}^{p-1}(\Phi(x))}\sqrt[p]{\omega} = \frac{b}{\Phi(\operatorname{Nr}_{L/K}(x))}\sqrt[p]{\omega} = \Phi(\zeta_p)\sqrt[p]{\omega} = \zeta_p^{-e^{p-2}}\sqrt[p]{\omega}.$$

Hence, $\tilde{\sigma}$ is of order p^2 and $\tilde{\lambda} = \tilde{\sigma}^p$. Finally, the relation $\tilde{\eta}\tilde{\sigma} = \tilde{\sigma}^{p+1}\tilde{\eta}$ holds. It follows that the group isomorphism $\varphi : \operatorname{Gal}(M/K) \cong C_{p^2} \rtimes C_p$, sending $\tilde{\sigma} \mapsto u$, $\tilde{\eta} \mapsto v$, is a solution isomorphism for the given embedding problem.

$$\frac{\bar{\tau}\omega}{\omega^e} = \frac{\bar{\tau}(\Phi(\beta))}{\Phi(\beta)^e} \cdot \frac{\bar{\tau}(\Phi(\sqrt[p]{a}))}{\Phi(\sqrt[p]{a})^e} = \left((\beta\sqrt[p]{a})^{\frac{1-e^{p-1}}{p}}\right)^p \in L^{*p} \quad , \frac{\bar{\tau}b}{b^e} = \left(\operatorname{Nr}_{L/K}(x)^{\frac{1-e^{p-1}}{p}}\right)^p \in L^{*p}.$$

Thus, $M/\mathbb{Q}$ is a Galois extension. We extend $\bar{\tau}$ to $\kappa \in \operatorname{Gal}(M/\mathbb{Q})$ by

$$\kappa(\sqrt[p]{\omega}) = (\sqrt[p]{a}\beta)^{(1-e^{p-1})/p}\left(\sqrt[p]{\omega}\right)^e, \qquad \kappa(\sqrt[p]{b}) = \operatorname{Nr}_{L/K}(x)^{(1-e^{p-1})/p}(\sqrt[p]{b})^e.$$

The rest remains exactly as in the previous case. □

We end this section with the following observations.

Theorem 4.3. *(a) Suppose that $L = \mathbb{Q}(\zeta_{p^2})$. Then $x \in L^*$ induces an H_{p^3}-extension if and only if it induces a $C_{p^2} \rtimes C_p$-extension (in the ways described in Theorem 4.1 and Theorem 4.2).*
(b) Suppose that $L \neq \mathbb{Q}(\zeta_{p^2})$. If an element $x \in L^*$ does not induce an H_{p^3}-extension, it necessarily induces a $C_{p^2} \rtimes C_p$-extension.
(c) There are infinitely many elements $x \in L^*$ which induce both an H_{p^3}-extension and a $C_{p^2} \rtimes C_p$-extension.

Proof. (a) and (b) follow from $\Phi(\zeta_p) = \zeta_p^{-e^{p-2}}$. (c) follows from Corollary 3.4. □

5. Polynomials for the non-abelian groups of order 27

Following Ledet, we shall construct E-extensions and their polynomials over $\mathbb{Q}$, induced by the elements x we found in Example 3.6 and Example 3.7. We replace the primitive root $e = 2$ with $e = -1$ - the homomorphism Φ does not depend on e when we consider it modulo 3 (it may not be the case for higher primes). Clearly, $\kappa\sqrt[3]{\omega} = 1/\sqrt[3]{\omega}$, thus $\bar{\tau}\omega = 1/\omega$, so $\omega + 1/\omega \in F$. Set $X = \sqrt[3]{\omega} + 1/\sqrt[3]{\omega}$. Then $X^3 - 3X = \omega + 1/\omega$. Hence, if $p(X) \in \mathbb{Q}[X]$ is the minimal polynomial for $\omega + 1/\omega$ over $\mathbb{Q}$, then the splitting field of $p(X^3 - 3X)$ over $\mathbb{Q}$ is $F(\sqrt[3]{\omega} + 1/\sqrt[3]{\omega})$, i.e, $p(X^3 - 3X)$ is an E-polynomial over $\mathbb{Q}$.

Example 5.1. Let x be as in Example 3.7. In order to get an H_{27}-polynomial over $\mathbb{Q}$ induced by this x, consider $\omega = \Phi(x^2\bar{\sigma}x)$. Using mathematical software (Maple), we get the following $\mathbb{H}_{27}$-polynomial over $\mathbb{Q}$:

$$p(X^3 - 3X) = (X^3 - 3X)^3 - \frac{3^4}{7^2}(X^3 - 3X)^2 - \frac{3 \cdot 37}{7^3}(X^3 - 3X) + \frac{1489}{7^4}.$$

Example 5.2. Due to the presence of the radical $\sqrt[p]{a}$, explicit extensions for the $C_{p^2} \rtimes C_p$ are more complicated to describe (unless $L = \mathbb{Q}(\zeta_{p^2})$).

Let x be as in Example 3.6. In order to get a $C_9 \rtimes C_3$-polynomial over $\mathbb{Q}$ induced by this x, consider $\omega = \Phi(x^2\bar{\sigma}x \cdot \sqrt[3]{a})$, where $L = K(\sqrt[3]{a})$. Denote $\delta = \delta_3(7)$ and consider the element

$$\theta = 3\delta^2 + 3\delta + 3\zeta_3\delta + \zeta_3 - 4.$$

The minimal polynomial for δ over $\mathbb{Q}$ is $X^3 + X^2 - 2X - 1$, and $\sigma\delta = \delta^2 - 2$. It follows that $\bar{\sigma}\theta = \zeta_3\theta$. Therefore, we can take $\sqrt[3]{a} = \theta$. Using mathematical software (Maple), we get the following $C_9 \rtimes C_3$-polynomial over $\mathbb{Q}$:

$$p(X^3-3X) = (X^3-3X)^3 - \frac{2\cdot 3^2\cdot 29}{13^2}(X^3-3X)^2 - \frac{3\cdot 5\cdot 373}{13^3}(X^3-3X) + \frac{6791}{13^3\cdot 7}.$$

6. ACKNOWLEDGMENT

The author is grateful to Jack Sonn, Moshe Roitman and Arne Ledet for useful discussions.

REFERENCES

[1] F.Demeyer, E.Ingraham, *Separable Algebras Over Commutative Rings*, Springer-Verlag, 1971.

[2] V.V.Ishkhanov, B.B.Lur'e & D.K.Faddeev, *The Embedding problem in Galois Theory*, Translations of Mathematical Monographs 165, American Mathematical Society, 1997.

[3] A.Ledet, *Generic and Explicit Realization of Small p-Groups*, Journal of Symbolic Computation, **30**, 859-865, 2000.

[4] A.Ledet, *Brauer Type Embedding Problems*, Fields Institute Monographs, **21**, American Mathematical Society, 2005.

[5] C.U.Jensen, A.Ledet, Noriko Yui, *Generic Polynomials, Constructive aspects of the Inverse Galois Problem*, Mathematical Sciences Research Institute Publications, **45**, 2002.

[6] D.A. Marcus, *Number Fields*, Springer-Verlag, 1977.

[7] R.Massy, *Galois Averages*, Journal of Number Theory, **113**, 244-275, 2005.

[8] P.Ribenboim, *Classical Theory of Algebraic Numbers*, Springer-Verlag, 2001.

[9] M.Suzuki, *Group Theory II*, Springer-Verlag, 1986.

[10] J.-P. Serre, *Topics in Galois Theory*, Jones and Bartlett, Boston, 1992.

Oz Ben-Shimol
Department of Mathematics
University of Haifa
Mount Carmel 31905, Haifa, Israel
E-mail Address: obenshim@math.haifa.ac.il